MycoKeys 51: 1–14 (2019)
doi: 10.3897/mycokeys.51.32272
http://mycokeys.pensoft.net

RESEARCH ARTICLE

A peer-reviewed open-access journal
MycoKeys
Launched to accelerate biodiversity research

# The genus *Castanediella*

Chuan-Gen Lin[1,2], Darbhe J. Bhat[3,4], Jian-Kui Liu[5], Kevin D. Hyde[2], Yong Wang[1]

**1** *Department of Plant Pathology, College of Agriculture, Guizhou University, Guiyang, Guizhou 550025, China* **2** *Center of Excellence in Fungal Research, Mae Fah Luang University, Chiang Rai 57100, Thailand* **3** *128/1-J, Azad Housing Society, Curca, Goa Velha 403108, India* **4** *Formerly, Department of Botany, Goa University, Goa, India* **5** *Center for Bioinformatics, School of Life Science and Technology, University of Electronic Science and Technology of China, Chengdu 611731, China*

Corresponding author: *Yong Wang* (yongwangbis@aliyun.com)

Academic editor: *C. Gueidan* | Received 9 December 2018 | Accepted 23 January 2019 | Published 15 April 2019

**Citation:** Lin C-G, Bhat DJ, Liu J-K, Hyde KD, Wang Y (2019) The genus *Castanediella*. MycoKeys 51: 1–14. https://doi.org/10.3897/mycokeys.51.32272

## Abstract

Two new species, *Castanediella brevis* and *C. monoseptata*, are described, illustrated and compared with other *Castanediella* taxa. Evidence for the new species is provided by morphological comparison and sequence data analyses. *Castanediella brevis* can be distinguished from other *Castanediella* species by the short hyaline conidiophores and fusiform, aseptate hyaline conidia, while *C. monoseptata* differs from other *Castanediella* species by its unbranched conidiophores and fusiform, curved, 0–1-sepatate, hyaline conidia. Phylogenetic analysis of combined ITS and LSU sequence data was carried out to determine the phylogenetic placement of the species. A synopsis of hitherto described *Castanediella* species is provided. In addition, *Castanediella* is also compared with morphologically similar-looking genera such as *Idriella*, *Idriellopsis*, *Microdochium*, *Neoidriella*, *Paraidriella* and *Selenodriella*.

**Keywords**

new taxa, Castanediellaceae, hyphomycetes, phylogeny, Sordariomycetes

## Introduction

Hernández-Restrepo et al. (2017) introduced the family *Castanediellaceae* for the genus *Castanediella* within Xylariales and it was consolidated in recent study by Wijayawardene et al. (2018). The asexual morphs in *Castanediellaceae* are hyphomycetous and characterized by macronematous, mononematous or sporodochial, branched, brown to pale brown conidiophores, with monoblastic or polyblastic, sympodial, discrete, cylindrical to lageni-

Copyright *Chuan-Gen Lin et al.* This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

form, hyaline to subhyaline conidiogenous cells, that produce unicellular or transversely septate, cylindrical, fusiform or lunate, hyaline conidia (Hernández-Restrepo et al. 2017).

The genus *Castanediella* was established by Crous et al. (2015) to accommodate *C. acaciae*, *C. cagnizarii* and *C. ramosa* within *Xylariales* genera *incertae sedis*. The genus contains twelve species (Costa et al. 2018; Wanasinghe et al. 2018), each characterized by branched, hyaline to pale brown conidiophores, holoblastic, sympodial conidiogenous cells and falcate, cylindrical or fusiform, 0–3-sepate, hyaline conidia (Crous et al. 2015; Costa et al. 2018).

During a survey of hyphomycetes in Thailand, two hyaline-spored hyphomycetes were collected. They were shown to belong to the genus *Castanediella* based on morphology and phylogeny analyses of ITS and LSU sequence data. The new species *C. brevis* and *C. monoseptata* are introduced.

## Materials and methods

### Collection and isolation of fungi

Dead leaves from a variety of plants in two forests (Lampang province and Chiang Mai province) were collected in 2016 in Thailand. Samples were taken to the laboratory in Zip-lock plastic bags for examination. The specimens were incubated in sterile moist chambers and examined using a Motic SMZ 168 series microscope. Fungi were removed with a needle and placed in a drop of distilled water on a slide for morphological study. Photomicrographs of fungal structures were captured with a Canon 600D digital camera attached to a Nikon ECLIPSE Ni compound microscope. All measurements were made by the Tarosoft (R) Image FrameWork program. Photo-plates were made with Adobe Photoshop CS3 (Adobe Systems, USA). Isolation of the fungi on to potato dextrose agar (PDA) was performed by the single spore isolation method (Chomnunti et al. 2014). Dried material was deposited in the Herbarium of Mae Fah Luang University (**MFLU**), Chiang Rai, Thailand and herbarium of Kunming Institute of Botany, Chinese Academy of Sciences (**HKAS**), Kunming, China. Cultures were deposited at Mae Fah Luang University Culture Collection (**MFLUCC**), Chiang Rai, Thailand and Kunming Institute of Botany, Chinese Academy of Sciences (**KUMCC**), Kunming, China. FacesofFungi and Index Fungorum numbers were registered (Jayasiri et al. 2015; Index Fungorum 2018).

### DNA extraction, PCR amplification and sequencing

Genomic DNA was extracted from fungal mycelium grown on PDA or malt extract agar (MEA) at room temperature using the Fungal gDNA Kit (BioMIGA, USA) according to the manufacturer's instructions. The internal transcribed spacer region of ribosomal DNA (ITS) and large subunit nuclear ribosomal DNA (LSU) genes were amplified via polymerase chain reaction (PCR) using the following primers: ITS5 and

ITS4 (White et al. 1990) for ITS, and LR0R and LR5 (Vilgalys and Hester 1990) for LSU. The PCR products were sequenced with the same primers. The PCR amplification was performed in a 25 μL reaction volume containing 12.5 μL of 2 × Power Taq PCR MasterMix (a premix and ready to use solution, including 0.1 Units/μl Taq DNA Polymerase, 500 μM dNTP Mixture each [dATP, dCTP, dGTP, dTTP], 20 mM Tris-HCl pH 8.3, 100 Mm KCl, 3 mM $MgCl_2$, stabilizer and enhancer), 1 μL of each primer (10 μM), 1 μL genomic DNA extract and 9.5 μL deionised water. The PCR thermal cycle program of ITS and LSU were followed as: initially 94 °C for 3 min., followed by 35 cycles of denaturation at 94 °C for 30 s, annealing at 55 °C for 50 s, elongation at 72 °C for 1 min., and final extension at 72 °C for 10 min.

## Phylogenetic analyses

Original sequences were checked using BioEdit version 7.0.5.3 (Hall 1999), and most reference sequences were originated from previous publications. The remaining homogenous sequences were obtained by BLAST searches (Altschul et al. 1990) from GenBank. All sequences used in this study are listed in Table 1. Alignments for each locus were done in MAFFT v7.307 online version (Katoh and Standley 2016) and manually verified in MEGA 6.06 (Tamura et al. 2013). After alignment, the concatenation of different genes was done in SequenceMatrix 1.8 (Vaidya et al. 2011). The interleaved NEXUS files for Bayesian inference analyses were formatted with AliView v1.19-beta1k (Larsson 2014). Maximum parsimony (MP), maximum likelihood (ML) and Bayesian inference (BI) were used for phylogenetic analyses.

The best models of evolution for each gene region were determined using Akaike information criterion (AIC) as implemented in MrModeltest v2 (Nylander 2004). The analyses' results showed that the models GTR+I and GTR+I+G were the best ones for LSU and ITS sequence data, respectively.

MP analyses were performed in PAUP*4.0b10 (Swofford 2002) following Liu et al. (2016).

ML analyses were carried out in raxmlGUI v 1.5b1 (Silvestro and Michalak 2012) with RAxML v8.2.10 (Stamatakis 2014), using the ML + rapid bootstrap setting and the GTRGAMMAI (viz., GTR + GAMMA + I) substitution model with 1000 bootstrap replicates.

For BI analysis, Posterior probabilities (PP) (Rannala and Yang 1996; Zhaxybayeva and Gogarten 2002) were determined by Markov Chain Monte Carlo sampling (BMCMC) in MrBayes v 3.2.6 (Ronquist et al. 2012). For the combined dataset, the models were set to nst = 6 and rates = propinv for LSU and nst = 6 and rates = invgamma for ITS. Two independent analyses of two parallel runs and six simultaneous Markov chains were run for 1,000,000 generations, trees were sampled every 100$^{th}$ generation and the temperature value of the heated chains was set at 0.15. The first 25% sampled trees of each run were discarded as "burn-in", and the remaining trees were used for calculating posterior probabilities (PP) in the majority rule consensus tree with the sumt command in MrBayes.

Phylogenetic trees were drawn with TreeView 1.6.6 (Page 1996).

**Table 1.** GenBank accession numbers of isolates included in this study.

| Taxa | Isolate[a] | ITS | LSU |
|---|---|---|---|
| *Castanediella acaciae* | CPC 24869, CBS 139896 | NR_137985 | KR476763 |
| *Castanediella brevis* | KUMCC 18-0132 | MH806361 | MH806358 |
| *Castanediella cagnizarii* | MUCL 41095 | KC775732 | KC775707 |
| *Castanediella cagnizarii* | CBS 101043 | KP859051 | KP858988 |
| *Castanediella cagnizarii* | CBS 542.96 | KP859054 | KP858991 |
| *Castanediella camelliae* | CNUFC-DLHBS5-1 | MF926620 | MF926614 |
| *Castanediella camelliae* | CNUFC-DLHBS5-2 | MF926621 | MF926615 |
| *Castanediella communis* | CPC 27631 | KY173393 | – |
| *Castanediella couratarii* | CBS 579.71 | NR_145250 | KP858987 |
| *Castanediella eucalypti* | CPC 24746, CBS 139897 | NR_137981 | KR476758 |
| *Castanediella eucalypticola* | CPC 26539 | NR_145254 | KX228317 |
| *Castanediella eucalyptigena* | CBS 143178, CPC 32055 | MG386036 | MG386089 |
| *Castanediella hyalopenicillata* | CPC 25873 | KX306751 | KX306780 |
| *Castanediella malaysiana* | CPC 24918 | NR_154810 | KX306781 |
| *Castanediella monoseptata* | KUMCC 18-0133 | MH806360 | MH806357 |
| *Castanediella ramosa* | MUCL 39857 | KC775736 | KC775711 |
| *Subsessila turbinata* | MFLUCC 15-0831 | KX762288 | KX762289 |

[a] **CBS**, Centraalbureau voor Schimmelcultures, Utrecht, Netherlands; **CPC**, Culture collection of Pedro Crous, housed at CBS; **KUMCC**, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, China; **MFLUCC**, Mae Fah Luang University Culture Collection, Chiang Rai, Thailand; **MUCL**, Mycothèque de l'Université Catholique de Louvian, Belgium.




**Figure 1.** Phylogenetic tree generated from MP analysis based on combined LSU and ITS sequence data for the genus *Castanediella*. Bootstrap support values for maximum parsimony (MP, first set) and maximum likelihood (ML, second set) greater than 50% are indicated above or below the nodes. Ex-type strains are in bold, the new isolates are in red. The tree is rooted with *Subsessila turbinata* (MFLUCC 15-0831).

## Results

### Molecular phylogeny

The aligned sequence matrix comprises LSU and ITS sequence data for 16 taxa (ingroup) and one outgroup taxon with a total of 1438 characters after alignment including the gaps, of which 120 were parsimony informative, 77 parsimony-uninformative, and 1241 characters constant. The dataset consists of thirteen species within the genus. The tree was rooted with *Subsessila turbinata* (MFLUCC 15-0831). Maximum parsimony analysis resulted in two trees with TL = 391, CI = 0.657, RI = 0.642, RC = 0.422, HI = 0.343. For the Bayesian analysis, two parallel runs with six chains were run for 1,000,000 generations and trees were sampled every 100$^{th}$ generation, resulting in 20002 trees from two runs of which 15002 trees were used to calculate the posterior probabilities (each run resulted in 10001 trees of which 7501 trees were sampled).The MP and ML (lnL = -4041.301739) analyses based on combined LSU and ITS sequence data provided similar tree topologies, and the result of MP analysis is shown in Fig. 1.

The novelty of the species, *Castanediella brevis* and *C. monoseptata*, described in this study are supported by sequence data analyses as belonging to the genus *Castanediella*, but with low bootstrap support values. Isolates of *Castanediella brevis* and *C. monoseptata* formed separate clades in the phylogenetic inference, respectively. *Castanediella brevis* is sister to *C. malaysiana* and *C. ramosa*, while *C. monoseptata* shows close phylogenetic relationship to *C. couratarii* and *C. malaysiana*. Both the new taxa can be recognized as phylogenetically distinct species and are clearly novel based on the recommendations for molecular data (Jeewon and Hyde 2016).

MP, ML and BI were used for phylogenetic analyses in this study. The tree topologies of MP and ML resulted from the combined LSU and ITS sequence data are similar, but most of the nodes are in low bootstrap support (Fig. 1). Polytomy structure was formed in the BI tree generated from the combined LSU and ITS sequence data. More sequence data, especially the protein-coding genes, e.g. TEF1-α, RPB2, β-tubulin, are required in the future study of the genus *Castanediella*.

### Taxonomy

***Castanediella brevis*** **C.G. Lin & K.D. Hyde, sp. nov.**
MycoBank number: MB828879
Facesoffungi number: FoF04929
Figure 2

**Holotype.** THAILAND. Lampang: Amphoe Mueang Pan, Tambon Chae Son, on decaying leaves, 24 September 2016, Chuangen Lin, LCG 10-1 (MFLU 18-1695, holotype; HKAS 102198, isotype), ex-type living cultures KUMCC 18-0132.

**GenBank number.** ITS: MH806361, LSU:MH806358

**Etymology.** In reference to the short conidiophores.




**Figure 2.** *Castanediella brevis* (MFLU 18-1695, holotype) **a** host material **b** conidiophores on the host surface **c–g** conidiophores, conidiogenous cells with conidia **h** conidia. Scale bars: 10 μm (**c–g**), 5 μm (**h**).

*Saprobic* on plant host. **Asexual morph:** *Colonies* on substrate effuse, white. *Mycelium* partly superficial, composed of septate, branched, smooth, hyaline to subhyaline hyphae. *Conidiophores* macronematous, mononematous, solitary, erect, unbranched,

straight or flexuous, short, 0–1-septate, hyaline, subcylindrical, ampulliform, smooth, often reduced to conidiogenous cells. *Conidiogenous cells* holoblastic, polyblastic, sympodial, integrated, terminal, subcylindrical, ampulliform, hyaline, denticulate, with 2–4 tiny protuberant denticles, 3–14 × 1.5–5.5 μm. *Conidia* solitary, dry, acropleurogenous, smooth, fusiform, curved, aseptate, hyaline, 12.5–21.7 × 1.2–3 μm (av. 16.95 × 2.2 μm, n = 60). **Sexual morph**: Undetermined.

*Culture characteristics*: Conidia germinating on PDA within 24 h. Colonies on PDA effuse, greyish white to dark from above and below, reaching a diam. of 5–7 cm in 30 days at 25 °C.

**Notes.** Based on a megablast search of the NCBI nucleotide database using the ITS sequence of the ex-type culture, the highest similarities found were with *Castanediella malaysiana* (GenBank NR_154810; identities = 526/537(98%), gaps = 1/537(0%)) and *C. couratarii* (GenBank KX960789; identities = 521/538(97%), gaps = 3/538(0%)). *Castanediella brevis* differs from these two species by its conidiophore morphology. *Castanediella couratarii* has pale brown conidiophores and longer conidiogenous cells (10.5–37 × 2–3.5 μm) whereas *C. malysiana* has pale brown and longer conidiophores (76–157 × 2.5–3 μm).

Among the species that produce more or less falcate and aseptate conidia, *Castanediella communis*, *C. eucalypti*, *C. eucalypticola* and *C. eucalyptigena* are most similar to *C. brevis*. However, *Castanediella brevis* differs from these species by its short, unbranched and 0–1-septate conidiophores.

***Castanediella monoseptata*** **C.G. Lin & K.D. Hyde, sp. nov.**
MycoBank number: MB828881
Facesoffungi number: FoF04930
Figure 3

**Holotype.** THAILAND. Chiang Mai: on decaying leaves, 24 August 2016, Chuangen Lin, MRC 3-1 (MFLU 18-1696, holotype; HKAS 102199, isotype), ex-type living cultures KUMCC 18-0133.

**GenBank number.** ITS: MH806360, LSU: MH806357

**Etymology.** In reference to the 0–1-septate conidia

*Saprobic* on plant host. **Asexual morph:** *Colonies* on substrate effuse, white. *Mycelium* partly superficial, composed of septate, branched, hyaline to subhyaline, smooth hyphae. *Conidiophores* macronematous, mononematous, solitary, erect, unbranched, straight or flexuous, septate, hyaline, subcylindrical, smooth, 8–29 × 2–4 μm. *Conidiogenous cells* polyblastic, integrated, sympodial, subcylindrical, hyaline, with several scars. *Conidia* solitary, dry, acropleurogenous, smooth, fusiform, curved, 0–1-sepatate, hyaline, 15.4–25.8 × 1.5–2.3 μm (av. 23.03 × 1.98 μm, n = 45). **Sexual morph**: Undetermined.

*Culture characteristics*: Conidia germinating on PDA within 24 h. Colonies on PDA effuse, grayish white to dark from above and below, reaching a diam. of 5–7 cm in 30 days at 25 °C.




**Figure 3.** *Castanediella monoseptata* (MFLU 18-1696, holotype) **a** host material **b** conidiophores on the host surface **c–f** conidiophores, conidiogenous cells with conidia **g–l** conidia. Scale bars: 10 μm (**c, d**), 5 μm (**e–l**).

**Notes.** A megablast search of the NCBI nucleotide database using the ITS sequence of the ex-type culture showed the highest similarities with uncultured Sordariales fungi (GenBank GQ268569; identities = 518/539(96%), gaps = 3/539(0%)) and *Castanediella couratarii* (GenBank KX960789; identities = 516/540(96%), gaps = 4/540(0%)).

Five *Castanediella* species, *C. cagnizarii*, *C. diversispora*, *C. hyalopenicillata*, *C. malaysiana* and *C. ramosa*, were reported to produce 1-septate conidia. *Castanediella monoseptata* can be distinguished from these species by its unbranched conidiophores and falcate and 15.4–25.8 × 1.5–2.3 μm conidia. *Castanediella monoseptata* is phylogeneti-

cally closely related to *C. couratarii* and *C. ramosa*, but differs from both species by its conidial morphology. *Castanediella couratarii* has shorter conidia (9.5–19 × 2–3 µm) are aseptate and *C. ramosa* has larger conidia (26–44 × 2–3 µm) that are 0–3-septate.

## Discussion

In this study, two new *Castanediella* species, *C. brevis* and *C. monoseptata*, were identified from decaying leaves in Thailand and a synopsis of hitherto described *Castanediella* species is provided (Table 2).

**Table 2.** Synopsis of *Castanediella* species.

| Taxa | Conidiophores | Conidiogenous cells | Conidia | | | |
|---|---|---|---|---|---|---|
| | | | Shape | Size (µm) | Septa | Colour |
| *C. acaciae* | Subcylindrical, medium brown, 40–80 × 2–3 µm. | Polyblastic, ampulliform, pale brown, 10–15 × 2–3 µm. | Falcate with subobtuse ends | (8–)10–11(–12) × 1.5(–2) | 0 | Hyaline |
| *C. brevis* | Subcylindrical, ampulliform, hyaline, often reduced to conidiogenous cells | Polyblastic, cylindrical, hyaline, 3–14 × 1.5–5.5 µm | Fusiform, curved | 12.5–21.7 × 1.2–3.0 | 0 | Hyaline |
| *C. cagnizarii* | Cylindrical, brown at the base, subhyaline towards the apex, up to 45 µm long. | Polyblastic, sympodial, subhyaline, 5–22 × 3–4 µm. | Cylindrical to fusiform, curved at the ends | Two sizes, 10–15 × 2 or 20–26 × 2 | | Hyaline |
| *C. camelliae* | Conidiophores reduced to conidiogenous cell. | Cylindrical, ampulliform, globose to subglobose, or irregularly-shaped, 5.5–20.5 × 2–4.5 µm. | Straight to slightly curved, sometimes swollen in the middle part | 18.5–51.5 × 1.6–2.5 | Septum indistinct | Hyaline |
| *C. communis* | Subcylindrical, medium brown, 20–60 × 3–4 µm. | Polyblastic, subcylindrical to ampulliform, pale brown, 10–35 × 2–4 µm. | Falcate with subobtuse ends | (13–)17–20(–22) × (2–)2.5(–3) | 0 | Hyaline |
| *C. couratarii* | Pale brown | Lageniform to cylindrical, hyaline to pale brown, 10.5–37 × 2–3.5 µm | Lunate | 9.5–19 × 2–3 | 0 | Hyaline |
| *C. diversispora* | Pale brown to brown | Polyblastic, sympodial, pale brown to brown, 4–9 × 2–3.5 µm. | Type i) cylindrical, slightly uncinate at the ends, straight | Type i) 11.5–16 × 2 | Type i) 1-septate | Hyaline |
| | | | Type ii) cylindrical to slightly subacerose, slightly uncinate at the apex, abruptly attenuated at the base, straight | Type ii) 19.5–25 × 1.5–2 | Type ii) 1-septate | |
| | | | Type iii) long filiform, obtuse or rounded at the apex attenuated at the base, straight or curved | Type iii) 28.5–47 × 1 | Type iii) 1–3-septate | |

| Taxa | Conidiophores | Conidiogenous cells | Conidia | | | |
|---|---|---|---|---|---|---|
| | | | Shape | Size (µm) | Septa | Colour |
| *C. eucalypti* | Subcylindrical, medium brown, 10–30 × 3–4 µm. | Polyblastic, subcylindrical to ampulliform, pale brown, 8–25 × 2.5–4 µm. | Falcate, slightly curved, widest in middle with subobtuse ends | (15–)18–21(–23) × 2–3 | 0 | Hyaline |
| *C. eucalypticola* | Subcylindrical, medium brown, 5–30 × 3–5 µm. | Polyblastic, subcylindrical to ampulliform or lanceolate, pale brown, 5–20 × 3–3.5 µm. | Falcate, straight to curved, widest in the middle, apex subobtusely rounded, base truncate, 0.5 µm diam | (15–)20–26(–30) × (2.5–)3 | 0 | Hyaline |
| *C. eucalyptigena* | Subcylindrical, hyaline, frequently reduced to conidiogenous loci on hyphae, up to 15 µm tall, 3–5 µm diam. | Polyblastic, hyaline, ampulliform or subcylindrical, 2–10 × 2–5 µm | Falcate, tapering to acute ends that are subobtusely rounded | (13–)18–24(–30) × 2(–2.5) | 0 | Hyaline |
| *C. hyalopenicillata* | Cylindrical, penicillate, mono-, bi-, and terverticillate, hyaline, 24–69 × 1.5–3 µm. | Mono- and polyblastic, short cylindrical, ampulliform, hyaline, 6.5–14 × 2–4 µm | Fusiform, base pointed, apex obtuse | 14–24 × 2–3 | 0–1 | Hyaline |
| *C. malaysiana* | Cylindrical, biverticillate, pale brown, 76–157 × 2.5–3 µm. | Polyblastic, cylindrical, subcylindrical, hyaline, 19–28 × 2.5–3.5 µm. | Fusiform, curved, apex acuminate, and base acuminate or slightly flattened | 18–30 × 2–3 | 0–1 | Hyaline |
| *C. monoseptata* | Subcylindrical, unbranched, hyaline, 8–29 × 2–4 µm | Polyblastic, cylindrical, hyaline | Fusiform, curved | 15.4–25.8 × 1.5–2.3 | 0–1 | Hyaline |
| *C. ramosa* | Cylindrical, penicillate, brown at the base, subhyaline at the apex, up to 70 µm long | Polyblastic, subhyaline, 10–20 x 2.5–3.5 µm | Falcate | 26–44 × 2.2–3 | (0–) 1 (–3) | Hyaline |

Presently, the genus *Castanediella* contains 14 species, and is shown to be diverse in its habitats. Most of *Castanediella* species have been collected from plant leaves. *Castanediella acaciae*, *C. camelliae*, *C. communis*, *C. eucalypti*, and *C. eucalypticola* were isolated from disease symptoms on different host plant leaves (Crous et al. 2015, 2016a, b; Wanasinghe et al. 2018) whereas *C. cagnizarii* is the only species found on decaying leaves submerged in a stream (Castañeda Ruiz et al. 2005). Some *Castanediella* species were reported from decaying leaves, such as *C. brevis*, *C. cagnizarii*, *C. diversispora*, *C. hyalopenicillata* and *C. monoseptata* (Castañeda Ruiz et al. 2005; Hernández-Restrepo et al. 2016b; Costa et al. 2018). *Castanediella couratarii* was reported from dead wood (Hernández-Restrepo et al. 2016a).

The genus *Castanediella* is morphologically similar to *Idriella*, *Idriellopsis*, *Microdochium*, *Neoidriella*, *Paraidriella*, *Selenodriella* (Seifert et al. 2011; Crous et al.

2015; Hernández-Restrepo et al. 2016a). However, these genera can be distinguished by the branching pattern of their conidiophores and conidial shape and septation (Hernández-Restrepo et al. 2016a). *Castanediella* differs from these genera by its branched conidiophores, ampulliform conidiogenous cells with scars instead of denticles, and filiform, 0–1-septate, straight to curved conidia (Crous et al. 2015). These similar-looking genera are phylogenetically distinct (Crous et al. 2015; Hernández-Restrepo et al. 2016a). A comparative synopsis of these genera is provided (Table 3).

**Table 3.** Synopsis of *Castanediella*-like genera.

| Genera | Conidiophores | Conidiogenous cells | Conidia | Chlamydospores |
|---|---|---|---|---|
| *Castanediella* | Branched, pale brown to brown at the base and subhyaline at the apex. | Sympodial, small denticles or scars, subhyaline. | 0–1-sepate, falcate, lunate, cylindrical or fusiform, hyaline | Not observed. |
| *Idriella* | Brown, mostly reduced to conidiogenous cells | Denticulate, sympodial | Aseptate, lunate, curved, hyaline | Brown, uni- or pluricellular. |
| *Idriellopsis* | Unbranched, brown at the base, almost hyaline at the apex, mostly reduced to conidiogenous cells | Terminal, denticulate | 0–1-septate, falcate, curved, hyaline | Not observed |
| *Microdochium* | More or less verticillate, reduced to conidiogenous cells, hyaline | Hyaline, sympodial or percurrent, sometimes denticulate | Aseptate or multiseptate, lunate, falcate, fusiform, filiform, obovoid or subpyriform, straight or curved, hyaline | Terminal or intercalary, solitary, in chains or grouped in clusters, brown. |
| *Neoidriella* | Mostly unbranched, pale brown, mostly reduced to conidiogenous cells | Sympodial, denticulate, terminal. | Aseptate, cylindrical to obovoid, hyaline | Intercalary or terminal, pale brown. |
| *Paraidriella* | Unbranched, pale brown, mostly reduced to conidiogenous cells. | Sympodial, denticulate, terminal. | Aseptate, cylindrical to oblong, hyaline | Not observed. |
| *Selenodriella* | Unbranched or verticillate, brown. | Sympodial, denticulate, terminal and intercalary. | Aseptate, falcate, hyaline | Not observed |

## Acknowledgements

We would like to thank Dr. Shaun Pennycook (Landcare Research Manaaki Whenua, New Zealand) for advising on the fungal names. The research is supported by the Thailand Research grants entitled The future of specialist fungi in a changing climate: baseline data for generalist and specialist fungi associated with ants, *Rhododendron* species and *Dracaena* species (grant no: DBG6080013), Impact of climate change on fungal diversity and biogeography in the Greater Mekong Subregion (grant no: RDG6130001). Y. Wang would like to thank the projects of the National Natural Science Foundation of China (No. 31560489, 31500451), Talent project of Guizhou science and technology cooperation platform ([2017]5788-5), Bijie Science and Technology Project ([2016]19) and Guizhou science and technology department international cooperation base project ([2018]5806). J.K. Liu would like to thank the National Natural Science Foundation of China (NSFC 31600032) and the Science and Technology Foundation of Guizhou Province (LH [2015]7061).

## References

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. Journal of Molecular Biology 215: 403–410.

Castañeda Ruiz RF, Heredia GP, Arias RM, Stadler M, Minter DW (2005) Two Hyphomycetes from submerged plant material of México. Mycotaxon 91: 333–338.

Chomnunti P, Hongsanan S, Aguirre-Hudson B, Tian Q, Persoh D, Dhami MK, Alias AS, Xu J, Liu X, Stadler M, Hyde KD (2014) The sooty moulds. Fungal Diversity 66: 1–36. https://doi.org/10.1007/s13225-014-0278-5

Costa PM, Barbosa MA, Da Silva GV, Sosa D, Pérez-Martinez S, Castañeda-Ruiz RF, Malosso E (2018) *Castanediella diversispora* sp. nov. from the Brazilian Atlantic Forest. Mycotaxon 133: 63–69.

Crous PW, Wingfield MJ, Burgess TI, Hardy GESJ, Crane C, Barrett S, Cano-Lira JF, Leroux JJ, Thangavel R, Guarro J, Stchigel AM, Martín MP, Alfredo DS, Barber PA, Barreto RW, Baseia IG, Cano-Canals J, Cheewangkoon R, Ferreira RJ, Gené J, Lechat C, Moreno G, Roets F, Shivas RG, Sousa JO, Tan YP, Wiederhold NP, Abell SE, Accioly T, Albizu JL, Alves JL, Antoniolli ZI, Aplin N, Araújo J, Arzanlou M, Bezerra JDP, Bouchara JP, Carlavilla JR, Castillo A, Castroagudín VL, Ceresini PC, Claridge GF, Coelho G, Coimbra VRM, Costa LA, da Cunha KC, da silva SS, Daniel R, de beer ZW, Dueñas M, Edwards J, Enwistle P, Fiuza PO, Fournier J, García D, Gibertoni TB, Giraud S, Guevara-Suarez M, Gusmão LFP, Haituk S, Heykoop M, Hirooka Y, Hofmann TA, Houbraken J, Hughes DP, Kautmanová I, Koppel O, Koukol O, Larsson E, Latha KPD, Lee DH, Lisboa DO, Lisboa WS, López-Villalba Á, Maciel JLN, Manimohan P, Manjón JL, Marincowitz S, Marney TS, Meijer M, Miller AN, Olariaga I, Paiva LM, Piepenbring M, Poveda-Molero JC, Raj KNA, Raja HA, Rougeron A, Salcedo I, Samadi R, Santos TAB, Scarlett K, Seifert KA, Shuttleworth LA, Silva GA, Silva M, Siqueira JPZ, Souza-Motta CM, Stephenson SL (2016a) Fungal Planet description sheets: 469–557. Persoonia - Molecular Phylogeny and Evolution of Fungi 37: 218–403.

Crous PW, Wingfield MJ, Guarro J, Hernández-Restrepo M, Sutton DA, Acharya K, Barber PA, Boekhout T, Dimitrov RA, Duenas M, Dutta AK, Gene J, Gouliamova DE, Groenewald M, Lombard L, Morozova OV, Sarkar J, Smith MT, Stchigel AM, Wiederhold NP, Alexandrova AV, Antelmi I, Armengol J, Barnes I, Cano-Lira JF, Castaneda Ruiz RF, Contu M, Courtecuisse PR, da Silveira AL, Decock CA, de Goes A, Edathodu J, Ercole E, Firmino AC, Fouriem A, Fournier J, Furtado EL, Geering ADW, Gershenzon J, Giraldo A, Gramaje D, Hammerbacher A, He XL, Haryadi D, Khemmuk W, Kovalenko AE, Krawczynski R, Laich F, Lechat C, Lopes UP, Madrid H, Malysheva EF, Marin-Felix Y, Martin MP, Mostert L, Nigro F, Pereira OL, Picillo B, Pinho DB, Popov ES, Pelaez CAR, Rooney-Latham S, Sandoval-Denis M, Shivas RG, Silva V, Stoilova-Disheval MM, Telleria MT, Ullah C, Unsickern SB, van der Merwe NA, Vizzini A, Wagner HG, Wong PTW, Wood AR, Groenewald JZ (2015) Fungal Planet description sheets: 320–370. Persoonia 34: 167–266. https://doi.org/10.3767/003158515x688433

Crous PW, Wingfield MJ, Richardson DM, Leroux JJ, Strasberg D, Edwards J, Roets F, Hubka V, Taylor PWJ, Heykoop M, Martín MP, Moreno G, Sutton DA, Wiederhold NP, Barnes CW, Carlavilla JR, Gené J, Giraldo A, Guarnaccia V, Guarro J, Hernández-Restrepo M, Kola, Ik

M, Manjón JL, Pascoe IG, Popov ES, Sandoval-Denis M, Woudenberg JHC, Acharya K, Alexandrova AV, Alvarado P, Barbosa RN, Baseia IG, Blanchette RA, Boekhout T, Burgess TI, Cano-Lira JF, moková A, Dimitrov RA, Dyakov MY, Dueñas M, Dutta AK, Esteve-Raventós F, Fedosova AG, Fournier J, Gamboa P, Gouliamova DE, Grebenc T, Groenewald M, Hanse B, Hardy GESJ, Held BW, Jurjevi, Kaewgrajang T, Latha KPD, Lombard L, Luangsa-ard JJ, Lysková P, Mallátová N, Manimohan P, Miller AN, Mirabolfathy M, Morozova OV, Obodai M, Oliveira NT, Ordóñez ME, Otto EC, Paloi S, Peterson SW, Phosri C, Roux J, Salazar WA, Sánchez A, Sarria GA, Shin HD, Silva BDB, Silva GA, Smith MT, Souza-Motta CM, Stchigel AM, Stoilova-Disheva MM, Sulzbacher MA, Telleria MT, Toapanta C, Traba JM, Valenzuela-Lopez N, Watling R, Groenewald JZ (2016b) Fungal Planet description sheets: 400–468. Persoonia - Molecular Phylogeny and Evolution of Fungi 36: 316–458.

Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Nucleic Acids Symposium Series 41: 95–98.

Hernández-Restrepo M, Gené J, Castañeda-Ruiz RF, Mena-Portales J, Crous PW, Guarro J (2017) Phylogeny of saprobic microfungi from Southern Europe. Studies in Mycology 86: 53–97. https://doi.org/10.1016/j.simyco.2017.05.002

Hernández-Restrepo M, Groenewald JZ, Crous PW (2016a) Taxonomic and phylogenetic re-evaluation of *Microdochium*, *Monographella* and *Idriella*. Persoonia – Molecular Phylogeny and Evolution of Fungi 36: 57–82.

Hernández-Restrepo M, Schumacher RK, Wingfield MJ, Ahmad I, Cai L, Duong TA, Edwards J, Gené J, Groenewald JZ, Jabeen S (2016b) Fungal Systematics and Evolution: FUSE 2. Sydowia 68: 193–230.

Jayasiri SC, Hyde KD, Ariyawansa HA, Bhat J, Buyck B, Cai L, Dai Y-C, Abd-Elsalam KA, Ertz D, Hidayat I, Jeewon R, Jones EBG, Bahkali AH, Karunarathna SC, Liu J-K, Luangsa-ard JJ, Lumbsch HT, Maharachchikumbura SSN, McKenzie EHC, Moncalvo J-M, Ghobad-Nejhad M, Nilsson H, Pang K-L, Pereira OL, Phillips AJL, Raspé O, Rollins AW, Romero AI, Etayo J, Selçuk F, Stephenson SL, Suetrong S, Taylor JE, Tsui CKM, Vizzini A, Abdel-Wahab MA, Wen T-C, Boonmee S, Dai DQ, Daranagama DA, Dissanayake AJ, Ekanayaka AH, Fryar SC, Hongsanan S, Jayawardena RS, Li W-J, Perera RH, Phookamsak R, Silva NI, Thambugala KM, Tian Q, Wijayawardene NN, Zhao R-L, Zhao Q, Kang J-C, Promputtha I (2015) The Faces of Fungi database: fungal names linked with morphology, phylogeny and human impacts. Fungal Diversity 74: 3–18. https://doi.org/10.1007/s13225-015-0351-8

Jeewon R, Hyde KD (2016) Establishing species boundaries and new taxa among fungi: recommendations to resolve taxonomic ambiguities. Mycosphere 7: 1669–1677. https://doi.org/10.5943/mycosphere/7/11/4

Katoh K, Standley DM (2016) A simple method to control over-alignment in the MAFFT multiple sequence alignment program. Bioinformatics 32: 1933–1942. https://doi.org/10.1093/bioinformatics/btw108

Larsson A (2014) AliView: a fast and lightweight alignment viewer and editor for large data sets. Bioinformatics 30: 3276–3278. https://doi.org/10.1093/bioinformatics/btu531

Liu JK, Yang J, Maharachchikumbura SSN, McKenzie EHC, Jones EBG, Hyde KD, Liu ZY (2016) Novel chaetosphaeriaceous hyphomycetes from aquatic habitats. Mycological Progress 15: 1157–1167.

Nylander J (2004) MrModeltest v2. Program distributed by the author. Evolutionary Biology Centre, Uppsala University.

Page RDM (1996) TreeView: an application to display phylogenetic trees on personal computers. Computer Applications in the Biosciences 12: 357–358. https://doi.org/10.1093/bioinformatics/12.4.357

Rannala B, Yang Z (1996) Probability distribution of molecular evolutionary trees: A new method of phylogenetic inference. Journal of Molecular Evolution 43: 304–311. https://doi.org/10.1007/BF02338839

Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. Systematic Biology 61: 539–542.

Seifert KA, Morgan-Jones G, Gams W, Kendrick B (2011) The genera of hyphomycetes. CBS-KNAW Fungal Biodiversity Centre, Utrecht, The Netherlands.

Silvestro D, Michalak I (2012) raxmlGUI: a graphical front-end for RAxML. Organisms Diversity & Evolution 12: 335–337.

Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics 30: 1312–1313. https://doi.org/10.1093/bioinformatics/btu033

Swofford DL (2002) PAUP*: Phylogenetic Analysis Using Parsimony and other methods, version 4.0 b10. Sinauer Associates.

Tamura K, Stecher G, Peterson D, Filipski A, Kumar S (2013) MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. Molecular Biology and Evolution 30: 2725–2729. https://doi.org/10.1093/molbev/mst197

Vaidya G, Lohman DJ, Meier R (2011) SequenceMatrix: concatenation software for the fast assembly of multi-gene datasets with character set and codon information. Cladistics 27: 171–180. https://doi.org/10.1111/j.1096-0031.2010.00329.x

Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. Journal of Bacteriology 172: 4238–4246.

Wanasinghe DN, Phukhamsakda C, Hyde KD, Jeewon R, Lee HB, Gareth Jones EB, Tibpromma S, Tennakoon DS, Dissanayake AJ, Jayasiri SC, Gafforov Y, Camporesi E, Bulgakov TS, Ekanayake AH, Perera RH, Samarakoon MC, Goonasekara ID, Mapook A, Li W-J, Senanayake IC, Li J, Norphanphoun C, Doilom M, Bahkali AH, Xu J, Mortimer PE, Tibell L, Tibell S, Karunarathna SC (2018) Fungal diversity notes 709–839: taxonomic and phylogenetic contributions to fungal taxa with an emphasis on fungi on Rosaceae. Fungal Diversity 89: 1–236. https://doi.org/10.1007/s13225-018-0395-7

White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (Eds) PCR protocols: a guide to methods and applications. Academic Press, California, 315–322.

Wijayawardene NN, Hyde KD, Lumbsch HT, Liu JK, Maharachchikumbura SSN, Ekanayaka AH, Tian Q, Phookamsak R (2018) Outline of Ascomycota: 2017. Fungal Diversity 88: 167–263. https://doi.org/10.1007/s13225-018-0394-8

Zhaxybayeva O, Gogarten JP (2002) Bootstrap, Bayesian probability and maximum likelihood mapping: exploring new tools for comparative genome analyses. BMC Genomics 3: 4–4. https://doi.org/10.1186/1471-2164-3-4